THE LANCET

Supplementary appendix

This appendix formed part of the original submission and has been peer reviewed. We post it as supplied by the authors.

Supplement to: Hansen CH, Michlmayr D, Gubbels SM, Mølbak K, Ethelberg S. Assessment of protection against reinfection with SARS-CoV-2 among 4 million PCR-tested individuals in Denmark in 2020: a population-level observational study. *Lancet* 2021; published online March 17. http://dx.doi.org/10.1016/S0140-6736(21)00575-4.

Supplementary materials

Contents

Table S1. Characteristics of the people who died before the second surge.	2
Table S2. Number of tests per person during the second surge, among people tested in the first surge	3
Table S3a. Model parameters for main analysis: Reinfection during the second surge.	4
Table S3b. Model parameters for the alternative cohort analysis	5
Figure S1. Time between testing	6
Figure S2. Kaplan-Meier curves of the time until (re)infection during follow-up	7

	Infected during first surge	Not infected during first surge
Total tested during first surge	11,727	521,044*
Total deaths before 1 Sep, 2020	659 (5.6%)	6773 (1.3)
Age distribution among those who died		
0-29 years	0 (0.0%)	32 (0.5%)
30-59 years	27 (4.1%)	520 (7.7%)
60-69 years	66 (10.0%)	934 (13.8%)
70-79 years	189 (28.7%)	2016 (29.8%)
80-89 years	250 (37.9%)	2232 (33.0%)
90+ years	127 (19.3%)	1039 (15.3%)

^{*}note 610 people who tested positive for the first time between the first and second surge of the epidemic have been removed.

Table S1. Characteristics of the people who died before the second surge among those tested in the first surge.

		Individuals	infected durin	g first surge			Individuals	not infected duri	ng first surge	
		Number of tests per person during second surge					Number of tests per person during second surge			
	Total	0	i	2-3	4+	Total	0	1	2-3	4+
Total	11,068	3525 (32)	2426 (22)	2719 (25)	2398 (22)	514,271	124165 (24)	117711 (23)	154359 (30)	118036 (23)
Positive test duri	ng second surge	ę								
Overall	72	0 (0)	7 (10)	26 (36)	39 (54)	16819	0 (0)	3019 (18)	7008 (42)	6792 (40)
By sex		` ′	` '	` '	` ′		. ,	` '	` '	, ,
Women	46	0(0)	4 (9)	13 (28)	29 (63)	10484	0 (0)	1586 (15)	4096 (39)	4802 (46)
Men	26	0 (0)	3 (12)	13 (50)	10 (38)	6335	0(0)	1433 (23)	2912 (46)	1990 (31)
By age		. ,	. ,	. ,	` /		. ,	` '	` '	, ,
0-19 years	4	0(0)	1 (25)	3 (75)	0(0)	1881	0 (0)	411 (22)	891 (47)	579 (31)
20-34 years	15	0 (0)	0 (0)	5 (33)	10 (67)	4789	0(0)	817 (17)	2058 (43)	1914 (40)
35-50 years	20	0 (0)	2 (10)	5 (25)	13 (65)	4358	0(0)	685 (16)	1744 (40)	1929 (44)
50-64 years	16	0 (0)	1 (6)	6 (38)	9 (56)	3925	0(0)	690 (18)	1567 (40)	1668 (42)
65-79 years	8	0 (0)	0(0)	4 (50)	4 (50)	1255	0 (0)	298 (24)	540 (43)	417 (33)
80+ years	9	0 (0)	3 (33)	3 (33)	3 (33)	611	0(0)	118 (19)	208 (34)	285 (47)
Time of first test	during second	surge	` '	` '	` ′		. ,	` '	` '	, ,
September	36	0 (0)	1 (3)	10 (28)	25 (69)	7196	0 (0)	271 (4)	2556 (36)	4369 (61)
October	13	0 (0)	1 (8)	5 (38)	7 (54)	3872	0 (0)	440 (11)	1805 (47)	1627 (42)
November	15	0 (0)	3 (20)	5 (33)	7 (47)	3180	0 (0)	802 (25)	1702 (54)	676 (21)
December	8	0 (0)	2 (25)	6 (75)	0 (0)	2571	0 (0)	1506 (59)	945 (37)	120 (5)
Time of last test of	during second s	urge								
September	4	0 (0)	1 (25)	2 (50)	1 (25)	447	0 (0)	271 (61)	164 (37)	12 (3)
October	5	0 (0)	1 (20)	4 (80)	0 (0)	1194	0 (0)	440 (37)	593 (50)	161 (13)
November	11	0 (0)	3 (27)	4 (36)	4 (36)	3390	0 (0)	802 (24)	1728 (51)	860 (25)
December	52	0 (0)	2 (4)	16 (31)	34 (65)	11788	0 (0)	1506 (13)	4523 (38)	5759 (49)

Note: Data are number of people with row percentages shown in parentheses.

Table S2. Number of tests per person during the second surge, among people tested in the first surge.

	Degrees of freedom	Estimate	Standard error	p-value	LR test for inclusion of variable
				•	
Intercept	1	-7.4840	0.0600	< 0.0001	
Exposed (i.e. positive test during surge 1)	1	-1.6339	0.1181	< 0.0001	
Female sex	1	-0.0959	0.0162	< 0.0001	
Age group 1: 0-5 years	1	-0.5501	0.0807		
Age group 2: 6-14 years	1	-0.0660	0.0648		
Age group 3: 15-24 years	1	0.3149	0.0561		
Age group 4: 25-34 years	1	0.1110	0.0556		
Age group 5: 35-44 years	1	0.0114	0.0559		
Age group 6: 45-54 years	1	0.0014	0.0556		
Age group 7: 55-64 years	1	-0.1355	0.0563		
Age group 8: 65-74 years	1	-0.4629	0.0620		
Age group 9: 75-84 years	1	-0.4086	0.0667		
Age group 10: 85+ years	0	ref	-	-	<0.0001, 9 d.
Test frequency group 1: 1-2 tests	1	-1.7539	0.0363		
Test frequency group 2: 3-5 tests	1	-0.4490	0.0299		
Test frequency group 3: 6-10 tests	1	-0.1755	0.0310		
Test frequency group 4: 11+	0	ref	-	-	<0.0001, 3 d.

Note: Model parameters were estimated in a Poisson regression: generalized linear model, with a Poisson distribution and log link.

Table S3a. Model parameters for main analysis: Reinfection during the second surge.

	Degrees of freedom	Estimate	Standard error	p-value	LR test for inclusion of variable
	1	0.0207	0.2520	-0.0001	
Intercept	l 1	-9.9297	0.2520	< 0.0001	
Exposed (i.e. positive test during surge 1)	1	-1.5512	0.0854	< 0.0001	
Male sex	1	0.0916	0.0088	< 0.0001	0.0001.0.10
Age group 1: 0-5 years	0	ref	- 0.0241		<0.0001, 9 d.f
Age group 2: 6-14 years	1	0.4871	0.0341		
Age group 3: 15-24 years	1	0.7826	0.0318		
Age group 4: 25-34 years	1	0.6261	0.0319		
Age group 5: 35-44 years	1	0.5043	0.0324		
Age group 6: 45-54 years	1	0.5211	0.0322		
Age group 7: 55-64 years	1	0.3778	0.0330		
Age group 8: 65-74 years	1	0.0791	0.0358		
Age group 9: 75-84 years	1	0.0849	0.0395		
Age group 10: 85+ years	1	0.4665	0.0470		
Test frequency group 1: 1-2 tests	0	ref	-		<0.0001, 3 d.f
Test frequency group 2: 3-5 tests	1	1.3447	0.0123		
Test frequency group 3: 6-10 tests	1	1.6399	0.0142		
Test frequency group 4: 11+	1	1.7994	0.0244		
Start month: October*	1	-1.0642	0.6292		
Start month: September	1	0.9981	0.2503		
Start month: August	1	0.7907	0.2502		
Start month: July	1	0.4565	0.2503		
Start month: June	1	0.1644	0.2502		
Start month: May	1	-0.0761	0.2502		
Start month: April	1	-0.2509	0.2503		
Start month: March	1	-0.4593	0.2526		
Start month: February	0	ref	_		<0.0001, 8 d.f

Notes: Model parameters were estimated in a Poisson regression: generalized linear model, with a Poisson distribution and log link. Start month indicates month of index test, i.e. the initial test that determines exposure status. *A few participants had their index test during the first two days of October, with the start of follow-up 90 days later at the very end of December.

Table S3b. Model parameters for the alternative cohort analysis: reinfection at least three months after first infection.




Figure S1. Time between testingDistribution of time from the first to repeat tests (showing data from 6,547,105 intervals contributed by 2,550,533 individuals).




Figure S2. Kaplan-Meier curves of the time until (re)infection during follow-up (log-rank p<0.0001).